



Letter to Editor

Defamation Against Healthcare Workers During COVID-19 Pandemic



Kazuki Shimizu^{1,2,3*}, Leesa Lin⁴

*Correspondence to: Kazuki Shimizu, Email: k.shimizu1@lse.ac.uk

Copyright: © 2022 The Author(s); Published by Kerman University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Shimizu K, Lin L. Defamation against healthcare workers during COVID-19 pandemic. *Int J Health Policy Manag.* 2022;11(5):720–721. doi:10.34172/ijhpm.2020.184

Received: 28 August 2020; Accepted: 20 September 2020; ePublished: 27 September 2020

Dear Editor,

While front-line healthcare workers (HCWs) combatting coronavirus disease 2019 (COVID-19) have commanded much respect and praise, they have been persecuted in Japan – a collective society where individual reputation and work ethic are traditionally viewed as important as one's life. HCWs, who were exposed to higher risk of infection due to lack of sufficient personal protective equipment and mentally exhausted in daily work, have another epidemic to fight: defamation. They have been targeted for discrimination and abuse in off-duty hours and stigmatized as "germs" in public spaces with many suffering social sanctions in forms of social exclusion and even bullying.²

Anxiety and fear among Japanese citizens were generated in January,3 and driven by the increase of nosocomial infections in February-April.4 Healthcare facilities and HCWs were seen as epicenters, which triggered widespread irrational prejudice and discrimination against HCWs. They were denied use of public buses, taxis, and even urged to vacate rental housing. Family members also suffered from defamation. Children of HCWs were asked to voluntarily refrain from using nursery schools or discriminated as "You are COVID-19" in their schools.2 Concurrently, there has been peer pressure among HCWs to commit to working on the front-line, fueled by the strong sense of responsibility that runs deep in Japanese work culture. HCWs had to struggle with the tension between the burnout and social sanctions, resulting in a wave of temporary or permanent leaves of absence² and consequently damaging the country's health system capacity.

To understand the discrimination against HCWs in Japan, root causes of the public anxiety towards COVID-19 must be investigated. First, public panic was incited by the interrupted access to testing.⁵ Due to the lack of testing capacity, only a

small portion of the infected cases has been identified,6-8 leading to public panic towards a largely underdiagnosed population. Unmet needs of health services are unprecedented and unfamiliar to Japanese residents who have historically enjoyed the resilient health system that ensured free access to healthcare. 9,10 Second, while designated medical institutions for infectious diseases continued accepting COVID-19 patients, some hospitals avoided them. In Tokyo, the number of rejection of ambulance cars amounted to 2365 in April 2020, which was nearly quadrupled compared to April 2019.11 Third, there has been a lack of governmental leadership to support HCWs; in fact, Japanese government's emergency risk communication was identified as a weakness that has negatively impacted its response to COVID-19.5 To date, there has been scant strong and clear messaging from political leaders that shows solidarity with front-line HCWs or any public acknowledgement of their commitment to protecting people's health during the pandemic. Fourth, there is a lack of strong medical journalism in Japan. Sensational media reports that are not based on scientific evidence have sometimes disseminated false information to the public.12 Fifth, Japan's declaration of State of Emergency, which was made between April and May, is not enforceable by law. All countermeasures such as remote working and staying indoors were on a voluntary basis. The strong peer pressure among Japanese has worked as a double-edged sword, which not only positively worked for infection control but negatively made Japanese residents suspicious of each other. Vigilantes have emerged, and coronavirus harassment for those who do not confirm has further divided the society.¹³ HCWs were socially excluded by an agitated public and irrationally criticized as the source of COVID-19.

Even before the COVID-19 pandemic, the pressure of deep-rooted conformity and social sanction that intersected with public health threats in Japanese society have long been documented. In 2011, stigmatized evacuees of the Fukushima Daiichi nuclear disaster were bullied at school, ¹⁴ highlighting an urgent need for context-specific public health education tailored to the Japanese sociocultural environment. During the 2009 influenza A (H1N1) pandemic, discrimination against front-line HCWs was acknowledged as a critical public health crisis. ¹⁵ It has been a decade since Japan was tested with a pandemic, yet not much progress has been made despite the evidence that the feeling of being supported

by the government with frequent communication and encouragement will motivate HCWs.15

To continue the fight against COVID-19, public leaders must ensure citizens' access to testing, take responsibility of HCWs' safety, show strong solidarity supporting HCWs, and prepare effective communication strategies tackling discrimination and social sanction against front-line HCWs.

Ethical issues

Not applicable.

Competing interests

Authors declare that they have no competing interests.

Authors' contributions

KS conceived the presented data, collected, and analyzed the data. KS drafted the initial manuscript. LL critically reviewed and revised the manuscript. Both authors have reviewed, commented on, and approved the final version of the manuscript.

Funding

KS receives research support in the United Kingdom by The Rotary Foundation '[grant number: GG1986485];' Japan Student Services Organization '[grant number: NM1910100304];' and the British Council Japan Association. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' affiliations

¹Department of Health Policy, London School of Economics and Political Science, London, UK. ²Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London, UK. ³Graduate School of Medicine, Hokkaido University, Sapporo, Japan. 4Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK.

References

- McCauley L, Hayes R. Taking responsibility for front-line health-care workers. Lancet Public Health. 2020;5(9):e461-e462. doi:10.1016/ s2468-2667(20)30179-1
- Around 30% of health-care workers feel "prejudice and discrimination" towards themselves and their surroundings (in Japanese). M3 website. https://www.m3.com/news/iryoishin/757309. Accessed August 27, 2020. Published April 26, 2020.
- Shimizu K. 2019-nCoV, fake news, and racism. Lancet. 2020; 395(10225):685-686. doi:10.1016/s0140-6736(20)30357-3

- Tani Y, Sawano T, Kawamoto A, Ozaki A, Tanimoto T. Nosocomial SARS-CoV-2 infections in Japan: a crosssectional newspaper database survey. Int J Health Policy Manag. 2020;9(10):461-463. doi:10.34172/ijhpm.2020.75
- Shimizu K, Wharton G, Sakamoto H, Mossialos E. Resurgence of covid-19 in Japan. BMJ. 2020;370:m3221. doi:10.1136/bmj.m3221
- Yamamoto K, Ozaki A, Senoo Y, et al. Underperformance of reversetranscriptase polymerase chain reaction in Japan and potential implications from Diamond Princess cruise ship and other countries during the ongoing COVID-19 pandemic. Int J Health Policy Manag. 2020; In Press. doi:10.34172/ijhpm.2020.109
- Omori R, Mizumoto K, Chowell G. Changes in testing rates could mask the novel coronavirus disease (COVID-19) growth rate. Int J Infect Dis. 2020;94:116-118. doi:10.1016/j.ijid.2020.04.021
- Omori R, Mizumoto K, Nishiura H. Ascertainment rate of novel coronavirus disease (COVID-19) in Japan. Int J Infect Dis. 2020; 96:673-675. doi:10.1016/j.ijid.2020.04.080
- Shibuya K, Hashimoto H, Ikegami N, et al. Future of Japan's system of good health at low cost with equity: beyond universal coverage. Lancet. 2011;378(9798):1265-1273. doi:10.1016/s0140-6736(11)61098-2
- 10. Sakamoto H, Rahman M, Nomura S, et al. Japan Health System Review. New Delhi: World Health Organization, Regional Office for South-East Asia; 2018. https://apps.who.int/iris/handle/10665/259941. Accessed August 27, 2020.
- 11. NHK (Japan Broadcasting Corporation). New monitoring indicators in Tokyo (in Japanese). https://www3.nhk.or.jp/news/special/coronavirus/number-tokyo/new-monitoring.html. Accessed September 16, 2020). Published June 30, 2020.
- World Health Organization. Strengthening health security by implementing the International Health Regulations (2005). Joint external evaluation of IHR core capacities of Japan. Mission report: 26 February - 2 March 2018. https://www.who.int/ihr/publications/ WHO-WHE-CPI-REP-2018.23/en/. Accessed August 27, 2020.
- 13. NHK World. Coronavirus Leads to Worrying Rise in Social Shaming. https://www3.nhk.or.jp/nhkworld/en/news/backstories/1083/. Accessed August 27, 2020. Published May 12, 2020.
- Sawano T, Nishikawa Y, Ozaki A, Leppold C, Tsubokura M. The Fukushima Daiichi Nuclear Power Plant accident and school bullying of affected children and adolescents: the need for continuous radiation. education. J Radiat Res. 2018;59(3):381-384. doi:10.1093/jrr/rry025
- 15. Imai H, Matsuishi K, Ito A, et al. Factors associated with motivation and hesitation to work among health professionals during a public crisis: a cross sectional study of hospital workers in Japan during the pandemic (H1N1) 2009. BMC Public Health. 2010;10:672. doi:10.1186/1471-2458-10-672