Article title: The Use of Evidence to Design an Essential Package of Health Services in Pakistan: A Review and Analysis of Prioritisation Decisions at Different Stages of the Appraisal Process

Journal name: International Journal of Health Policy and Management (IJHPM)

Authors' information: Sergio Torres-Rueda1, Anna Vassall¹*, Raza Zaidi², Nichola Kitson¹, Muhammad Khalid², Wahaj Zulfiqar², Maarten

Jansen³, Wajeeha Raza⁴, Maryam Huda⁵, Frank Sandmann⁶, Rob Baltussen³, Sameen Siddiqi⁵, Ala Alwan⁷

¹Department of Global Health & Development, London School of Hygiene and Tropical Medicine, London, UK.

²Ministry of National Health Services, Regulations and Coordination, Islamabad, Pakistan.

³Department of Health Evidence, Radboud Institute of Health Sciences, Radboud University Medical Center, Nijmegen, The Netherlands.

⁴Centre for Health Economics, University of York, York, UK.

⁵Department of Community Health Sciences, Aga Khan University, Karachi, Pakistan.

⁶Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK.

⁷DCP3 Country Translation, London School of Hygiene and Tropical Medicine, London, UK.

*Correspondence to: Anna Vassall; Email: <u>Anna.Vassall@lshtm.ac.uk</u>

Citation: Torres-Rueda S, Vassall A, Zaidi R, et al. The use of evidence to design an essential package of health services in Pakistan: a review and analysis of prioritisation decisions at different stages of the appraisal process. Int J Health Policy Manag. 2024;13:8043. doi:10.34172/ijhpm.2024.8043

Supplementary file 1. Values Used for Each Intervention for Each Decision Criteria (Evidence and Evidence Quality)

Abbreviations: DCP, Disease Control Priorities; EUHC, Essential Universal Health Coverage; TWG, technical working group; NAC, National Advisory Council; EPHS, Essential Package of Health Services; IIP, immediate implementation package; World Health Organization; HiB, Haemophilus influenzae type b; BCG, Bacillus Calmette-Guéri; IRS, indoor residual spraying; HPV, human papillomavirus; STIs, sexually transmitted infections; TB, tuberculosis; WASH, water, sanitation and hygiene; G6PD, glucose-6-phosphate dehydrogenase; BEmONC, Basic Emergency Obstetric and Newborn Care; PLHIV, people living with HIV; IMAI, integrated management of adolescent and adult illness; CVD, cardiovascular disease; COPD, chronic obstructive pulmonary disease; ACEi, angiotensin-converting enzyme inhibitor; ARBs, angiotensin receptor blockers; IV, intravenous; CEmONC, Comprehensive Emergency Obstetric and Newborn Care; ART, antiretroviral treatment; NTDs, neglected tropical diseases; ECD, early childhood development; ARV, antiretroviral; IMCI, Integrated management of childhood illness; IUCDs, IDU, intravenous drug user; MSM, men who have sex with men; MDR, multidrug-resistant tuberculosis; XDR, extensively drug-resistant tuberculosis; C, community; HC, health centre; FLH, first-level hospital; RH, referral hosptal; ICER, incremental cost-effectiveness ratio

| DCP3 Code | Intervention name | Cost-effectiveness | Budget impact | Burden of disease | ICER quality | Current coverage |
|--------------|--|--------------------|---------------|----------------------|--------------|------------------|
| C1 | Antenatal and postpartum education on birth spacing | High | Low | Medium | Medium | High |
| C10 | Education on handwashing, personal hygiene and safe disposal of children's stool | High | Low | High | Medium | High |
| C11 | Pneumococcus vaccination | Medium | Medium | High | High | High |
| C12 | Rotavirus vaccination | Medium | Low | High | High | High |
| C14 | Vitamin A and zinc for children | Medium | High | High | Low | High |
| C16 | Childhood vaccination series (diphtheria, pertussis, tetanus, polio, BCG, measles, hepatitis B, HiB, rubella) | High | Medium | High | Medium | High |
| C17 | Indoor residual spraying | Medium | Medium | High | Medium | Low |
| C18 | Education of schoolchildren on oral health | Medium | High | Medium | Low | Low |
| C19 | Vision pre-screening by teachers; vision tests and provision of ready-made glasses on-site by eye specialists | Medium | High | High | Low | Low |
| C2 | Counselling of mothers on providing thermal care for pre- term new-borns (delayed bath and skin to skin contact) | High | Low | Medium | Medium | Low |
| C20 | School based HPV vaccination for girls | Low | Low | Low | High | No |
| C21 | Mass drug administration (NTDs) | Low | High | Medium | Low | No |
| C23 | Adolescent-friendly services for STIs | Medium | Low | Medium | Low | Low |
| C24 | Life skills training in schools | Low | High | High | Low | Low |
| C27a | Provision of iron and folic acid supplementation to pregnant women, and provision of food or caloric supplementation to pregnant women in food-insecure households | High | High | High | Medium | High |
| C27b | Provision of iron and folic acid supplementation to pregnant women, and provision of food or caloric supplementation to pregnant women in food-insecurity households | High | High | High | Medium | High |
| | Community-based HIV testing and counselling (for example, mobile units and venue- | | | | | |
| C28 | based testing), with appropriate referral or linkage to care and | High | Low | Low | Medium | Low |
| C30a | Provision of condoms to key populations, including sex workers, men who have sex with men, people who inject drugs, transgender populations, and prisoners | High | Low | Medium | Medium | Low |
| C30b | Provision of Disposable syringes who inject drugs (IDU) | High | Low | Medium | Medium | Low |

| DCP3 Code | Intervention name | Cost-effectiveness | Budget impact | Burden of disease | ICER quality | Current coverage |
|--------------|---|--------------------|---------------|----------------------|--------------|------------------|
| C32 | Routine contact tracing to identify individuals exposed to TB and link them to care | Low | Low | Low | Low | High |
| C33 | Test for G6PD deficiency | Low | Low | Low | Low | Medium |
| C3a | Management of labour and delivery in low-risk women by skilled attendant (CL) | High | Low | High | Low | Low |
| C3b | Basic neonatal resuscitation following delivery (CL) | High | Low | High | Low | Low |
| C3c | Management of labour and delivery in low-risk women by skilled attendant (PHC) | High | Low | High | Low | High |
| C3d | Basic neonatal resuscitation following delivery (PHC) | High | Low | High | Low | High |
| C34 | Environmental management for malaria | High | Medium | High | Low | Low |
| C4 | Promotion of breastfeeding and complementary feeding by community health workers | High | Low | High | Medium | High |
| C41 | Mass drug administration (malaria) | High | High | High | Low | No |
| C43 | Early detection and treatment of leishmaniasis, dengue, chikungunya, rabies, trachoma and helminthiasis. | Medium | Low | Medium | Low | Medium |
| C45 | Identify and refer patients with high risk | None | None | None | None | High |
| | In the context of an emerging infectious outbreak, provide advice and guidance on how to | | | | | |
| C46 | recognize early symptoms and signs and when to seek medical attention | Medium | Low | Medium | Low | High |
| C47 | Exercise-based pulmonary rehabilitation | Medium | Low | Low | Low | No |
| C48 | Self-managed treatment of migraine | Low | Low | Medium | Low | No |
| C5 | Tetanus toxoid immunization among schoolchildren and women attending antenatal care | High | Low | Medium | Medium | High |
| | Parent training of high-risk families, including nurse home visitation for child | | | | | |
| C50 | maltreatment | Medium | High | High | Low | Low |
| C51 | WASH behaviour change interventions, such as community led total sanitation | Low | Medium | Medium | Low | High |
| | Identification/screening of the early childhood development issues motor, sensory and | | | | | |
| C53a | language stimulation | Medium | High | High | Low | Low |
| C53b | ECD rehabilitation interventions | Medium | High | High | Low | Low |
| C56 | Interventions for wheelchair users | None | None | None | None | No |
| C8 | Acute severe malnutrition management | Low | High | High | Low | High |
| C9 | Integrated community case management | Medium | Low | Low | Low | Low |
| FLH1 | Care for foetal growth restriction | Medium | High | High | Low | Medium |
| | Surgical termination of pregnancy by maternal vacuum aspiration and dilatation & | | | | | |
| FLH10 | curettage | Low | Low | Low | Low | Low |
| FLH11 | Care for severe childhood infections | Medium | High | High | Low | Medium |
| FLH12 | Severe acute malnutrition management | Medium | Low | Low | Medium | Medium |
| FLH13 | Early detection and treatment of early-stage cervical cancer | Medium | Low | Low | Medium | Low |
| FLH14 | Insertion and removal of contraceptives | Medium | Low | Medium | Low | Low |
| FLH15 | Tubal ligation | Low | High | High | Low | Low |
| FLH16 | Vasectomy | Medium | Low | Medium | Low | Low |
| FLH17 | Referral of cases of treatment failure for drug susceptibility testing; enrolment of those with MDR-TB for treatment per WHO guidelines (either short- or long-term regimen) | High | Low | High | Low | High |
| | Evaluation and management of fever in clinically unstable individuals using WHO IMAI guidelines, including empiric parenteral antimicrobials and antimalarial and resuscitative | | | | | |
| FLH18 | measures for septic shock | High | Low | High | Medium | Medium |
| FLH2 | Induction of labour post-term | Medium | Low | Medium | Low | Medium |
| FLH20 | Management of acute coronary syndromes | Low | High | Medium | Low | Medium |

| DCP3 Code | Intervention name | Cost-effectiveness | Budget impact | Burden of disease | ICER quality | Current coverage |
|--------------|--|--------------------|---------------|----------------------|--------------|------------------|
| | Management of acute exacerbations of asthma and COPD using systemic steroids, inhaled | | | | | |
| FLH22 | beta-agonists and if indicated oral antibiotics and oxygen therapy | Low | High | Medium | Medium | Low |
| FLH23 | Medical management of acute heart failure | Medium | Low | High | Medium | Medium |
| FLH24 | Bowel obstruction management | High | Low | Medium | Low | Medium |
| FLH25 | Calcium and vitamin D supplementation for secondary prevention of osteoporosisE264 | Low | High | High | Low | Low |
| FLH26 | Combination therapy, including low-dose corticosteroids and generic disease-modifying antirheumatic drugs (including methotrexate), for individuals with moderate to severe rheumatoid arthritis | High | Low | Medium | Low | Medium |
| | In settings where sickle cell disease is a public health concern, universal newborn | | 2011 | cului | 2011 | |
| FLH27 | screening followed by standard prophylaxis against bacterial infections and malaria | Medium | Low | Low | Medium | Low |
| | In setting where specific single-gene disorders are a public health concern (for example thalassemia), retrospective identification of carriers plus prospective (premarital) | | - | | | |
| FLH28 | screening and counselling to reduce rates of conception | Medium | Low | Low | Low | Low |
| FLH3 | Jaundice management with phototherapy | Medium | Low | Low | Low | Medium |
| FLH30 | Intoxication/poisoning management | Medium | Low | Low | Low | Low |
| FLH31 | Appendectomy | Low | Medium | Medium | Medium | High |
| FLH32 | Assisted vaginal delivery using vacuum extraction or forceps | Medium | Low | Medium | Medium | Low |
| FLH33 | Craniotomy for trauma | Medium | Low | Low | Medium | Low |
| FLH34 | Colostomy for acute bowel obstruction/volvulus and injuries. | High | Low | Medium | Low | Medium |
| FLH35 | Escharotomy or fasciotomy | Medium | Low | High | Medium | Low |
| FLH36 | Management of non-displaced fractures | High | Low | High | Medium | Medium |
| FLH37 | Hernia Repair | Medium | Low | Medium | Medium | Medium |
| FLH38 | Hysterectomy for uterine rupture or intractable postpartum haemorrhage | High | Low | Medium | Medium | High |
| FLH39 | Irrigation and debridement of open fractures | Medium | Low | Medium | Low | Medium |
| FLH4 | Eclampsia management with magnesium sulphate, including initial stabilization at health centres | Medium | Low | High | Medium | High |
| FLH40 | Management of osteomyelitis, including surgical debridement | Medium | Medium | High | Low | Medium |
| FLH41a | Management of Septic Arthritis | Medium | Low | Low | Medium | Low |
| FLH41b | Placement of External Fixation and Use of Traction for Fractures | Medium | Low | Low | Medium | Low |
| FLH42 | Relief of urinary obstruction by catheterization for fractures | High | Low | High | Low | High |
| FLH43 | Removal of gallbladder, including emergency surgery | Medium | Low | Medium | Low | Medium |
| FLH44 | Repair of perforations (for example perforated peptic ulcer, typhoid ileal perforation) | High | Low | High | Low | High |
| FLH45 | Resuscitation with advanced measures | Medium | Low | Medium | Low | Low |
| FLH46 | Basic Skin grafting | Medium | Low | Low | Medium | Low |
| FLH48a | Trauma laparotomy | High | Low | High | Medium | High |
| FLH49 | Trauma-related amputations | High | Low | High | Low | Medium |
| FLH5 | Maternal sepsis management | Medium | Low | Medium | Medium | Medium |
| FLH50 | Tube thoracostomy | Medium | Low | Low | Low | Medium |
| FLH52 | Compression therapy for amputations, burns, and vascular or lymphatic disorders | Medium | Low | Low | Medium | Medium |
| FLH53 | Evaluation and acute management of swallowing dysfunction | Medium | Low | Low | Low | Medium |
| FLH57 | Prevention and relief of refractory suffering and acute pain related to surgery, serious injury or other serious, complex or life-limiting health problems | None | None | None | None | Low |

| DCP3 Code | Intervention name | Cost-effectiveness | Budget impact | Burden of disease | ICER quality | Current coverage |
|--------------|---|--------------------|---------------|----------------------|--------------|------------------|
| FLH58 | First level hospital pathology services | None | None | None | None | Medium |
| | Management of new-born complications infections, meningitis, septicaemia, pneumonia and other very serious infections requiring continuous supportive care (such as IV fluids | | | | | |
| FLH6 | and oxygen) | High | Low | High | Medium | Medium |
| FLH7 | Preterm labour management | Low | Medium | Low | Medium | Medium |
| FLH8 | Management of labour and delivery in high-risk women, including operative delivery (CEmONC) | Low | High | High | Low | Medium |
| FLH9 | Surgery for ectopic pregnancy | High | Low | Medium | Medium | Medium |
| HC1 | Early detection and treatment of neonatal pneumonia with oral antibiotics | High | Low | High | Medium | High |
| HC10 | Screening and management of diabetes (gestational diabetes or pre-existing type II diabetes) | Low | Low | Low | Medium | Low |
| HC11 | Management of labour and delivery in low-risk women (BEmONC), including initial treatment of obstetric or delivery complications prior to transfer (Also included in Surgery package of services) | High | High | High | Low | High |
| HC11 HC12 | | | | | | |
| HCIZ | Detection and treatment of childhood infections with danger signs (IMCI) Among all individuals who are known to be HIV+, immediate ART initiation with regular | High | High | High | Low | Low |
| HC13 | monitoring of viral load for | Low | Low | Low | Medium | Low |
| HC13 HC14 | Psychological treatment | Low | High | Medium | Medium | Low |
| HC14 HC16 | Post gender-based violence care | Low | Medium | Medium | Low | Low |
| псто | Syndromic management of common sexual and reproductive tract infections (for example | LOW | wedium | Medium | LOW | LOW |
| HC17 | urethral discharge, genital ulcer and others) | Medium | High | High | Low | High |
| HC19 | For individuals testing positive for hepatitis B and C, assessment of treatment eligibility by trained providers followed by initiation and monitoring of ART when indicated | Low | High | High | Medium | Low |
| HC2 | Miscarriage and abortions management | Low | Low | Medium | Low | Medium |
| HC20 | Hepatitis B and C testing of high-risk individuals identified in the national testing policy with appropriate referral of positive individuals to trained providers | Medium | Medium | Medium | Low | Low |
| HC21 | Partner notification and expedited treatment for common STIs including HIV | High | Medium | High | Medium | Medium |
| | Provider-initiated testing and counselling for HIV, STIs and hepatitis for all in contact with the health system in high- prevalence setting, including prenatal care with appropriate referral/ linkages to care including immediate ART initiation for those testing positive for | | | | | |
| HC23 | HIV | Medium | High | High | Medium | Medium |
| | Hepatitis B vaccination for high-risk populations, including healthcare workers, IDU, MSM, | | | | | |
| HC24 | household contacts and partners with multiple sex partners | Medium | Low | Low | Medium | Low |
| HC25 | Medical male circumcision | Medium | Medium | Medium | Low | Low |
| | For PLHIV and children under five who are close contacts or household members of individuals with active TB, perform symptom screening and chest radiograph; if there is no | | | | | |
| HC26 | active TB, provide isoniazid preventive therapy according to current WHO guidelines | High | Medium | High | Medium | Low |
| HC27 | Diagnosis of TB and first-line treatment | High | Low | High | Low | High |
| | Screening for HIV in all individuals with a diagnosis of active TB; if HIV infection is present, | <u> </u> | | 0 | | J |
| HC28 | start (or refer for) ARV treatment and HIV care | High | Low | Medium | Low | Low |
| HC29 | Latent-TB screening and IPT for PLHIV | High | Low | Medium | Low | Low |
| HC3 | Management of premature rupture of membranes, including administration of antibiotics | Low | Low | Low | Low | Low |

| DCP3 Code | Intervention name | Cost-effectiveness | Budget impact | Burden of disease | ICER quality | Current coverage |
|---------------|---|--------------------|---------------|----------------------|--------------|------------------|
| HC30 | Fever management for clinically stable | Medium | Low | Low | Low | High |
| HC32 | Provision of insecticide nets to U5 children and pregnant women attending health centres | High | Low | Medium | Medium | Medium |
| HC33 | Identify and refer for progressive illness ** | None | None | None | None | Medium |
| HC36 | Long-term combination therapy for persons with multiple CVD risk factors, including screening for CVD in community setting using non-lab-based tools to assess overall CVD risk | Low | Medium | Medium | Low | Low |
| | Low-dose inhaled corticosteroids and bronchodilators for asthma and for selected | | | | | |
| HC37 | patients with COPD | Low | Low | Low | Medium | Low |
| HC38 | Provision of aspirin for all cases of suspected acute myocardial infarction | Medium | Low | Low | High | Low |
| HC39a | Screening and ACEi or ARBs for kidney disease | Low | Low | Low | Low | Low |
| HC41 | Secondary prophylaxis for rheumatic fever | Low | Low | Low | Low | Low |
| HC42 | Treatment of acute pharyngitis for rheumatic fever | High | Low | Medium | Low | High |
| HC45 | Opportunistic screening for hypertension | Medium | High | High | Medium | Low |
| HC46 | Tobacco cessation counselling | High | High | High | Medium | No |
| HC48 | Support for caregivers of dementia patients | Medium | Low | Low | Medium | No |
| HC49 | Bipolar disorder management | Low | Medium | Low | Low | No |
| HC4a | Provision of condoms and hormonal contraceptives, including emergency contraceptives | High | Low | High | Medium | Low |
| | Provision of condoms and hormonal contraceptives, including insertion and removal of | | | | | |
| HC4b | contraceptives (PHC) | High | Low | High | Medium | Low |
| | Management of depression and anxiety disorders with psychological and generic | | | | | |
| HC50 | antidepressants therapy | Low | High | Medium | Low | Low |
| HC53 | Screening and brief alcohol intervention | Low | Low | Low | Low | No |
| HC55 | Primary prevention of osteoporosis | Low | Low | Medium | Low | Medium |
| HC56 | Screening for congenital hearing loss | Low | Medium | Medium | Medium | Low |
| HC57a | Dental extraction (PHC) | Medium | High | High | Low | Medium |
| HC57b | Dental extraction (FLH) | Medium | High | High | Low | Medium |
| HC58a | Drainage of dental abscess (PHC) | Low | High | Medium | Low | Low |
| HC59 | Drainage of superficial abscess | Low | Medium | Medium | Low | Medium |
| HC5a | Counselling on kangaroo care for new-borns (CL) | Medium | Low | Low | Low | Low |
| HC5b | Counselling on kangaroo care for new-borns (PHC) | Medium | Low | Low | Low | Low |
| HC6 | Management of neonatal sepsis, pneumonia and meningitis using injectable and oral antibiotics | High | Low | Low | Medium | Low |
| | | | | Low | Low | Low |
| HC60 HC61 | Non-displaced fractures management | Low None | Low None | None | None | Low |
| HC61 HC62 | Resuscitation with basic life support measures | | Low | Low | Low | Medium |
| | Suturing laceration | Low | - | - | - | |
| HC63a HC64 | Treatment of caries (PHC) Basic management of MNIs and disorders | Low | High Low | Medium Low | Low | Low |
| HC64 HC66 | Psychosocial support and counselling | Low | Low | LOW | Low | Low |
| 100 | , | LUW | LOW | LOW | LOW | LUW |
| HC67 | Expanded palliative care and pain control measures, including prevention and relief of all physical and psychological symptoms of suffering | None | None | None | None | Low |
| HC68 | Health centre pathology services ** | None | None | None | None | Low |
| HC7 | Pharmacological termination of pregnancy | High | Low | High | Low | Low |

| DCP3 Code | Intervention name | Cost-effectiveness | Budget impact | Burden of disease | ICER quality | Current coverage |
|--------------|--|--------------------|---------------|----------------------|--------------|------------------|
| HC9a | Screening of hypertensive disorders in pregnancy | Low | Low | Low | Low | High |
| HC9b | Screening and management of hypertensive disorders in pregnancy | Low | Low | Low | Low | High |
| | Systematic identification of individuals with TB symptoms among high-risk groups and | | | | | |
| P5 | linkages to care (active case finding) | Low | Medium | Medium | Medium | High |
| RH1 | Full supportive care for preterm new-borns | High | Low | High | Medium | Medium |
| | Specialized TB services, including management of MDR- and XDR-TB treatment failure and | | | | | |
| RH2 | surgery for TB | Medium | Low | Medium | Low | Medium |
| | Management of refractory febrile illness including etiologic diagnosis at reference | | | | | |
| RH3 | microbial laboratory | Medium | Low | Low | Low | Medium |
| RH4 | Management of acute ventilator failure due to acute exacerbations of asthma and COPD | Low | Low | Low | Medium | Low |
| | Retinopathy screening via telemedicine, followed by treatment using laser | | | | | |
| RH5 | photocoagulation | Medium | Low | Low | Medium | Low |
| | Use of percutaneous coronary intervention for acute myocardial infarction where | | | | | |
| RH6 | resources permit | Low | Medium | High | Low | Low |
| | Treatment of early-stage breast cancer with appropriate multimodal approaches | | | | | |
| | (including generic chemotherapy) with curative intent for cases detected by clinical | | | | | |
| RH7 | examination | Low | Low | Low | Medium | Medium |
| | Treatment of early-stage colorectal cancer with appropriate multimodal approaches | | | | | |
| DUIO | (including generic chemotherapy) with curative intent for cases detected by clinical | 1 | | 1 | No. dia ma | 1 |
| RH8 | examination | Low | Low | Low | Medium | Low |
| | Treatment of early-stage childhood cancers (such as Burkitt and Hodgkin lymphoma, | | | | | |
| RH9 | acute lymphoblastic leukaemia, retinoblastoma and Wilms tumour) with curative intent in paediatric cancer units or hospitals | 1 | 1 | Medium | Medium | 1.0 |
| КПЭ | Elective surgical repair of common orthopaedic injuries (for example meniscal and | Low | Low | wealum | wedium | Low |
| RH10 | ligamentous tears) in individuals with severe functional limitation | High | Medium | High | Medium | Low |
| KUIU | Urgent, definitive surgical management of orthopaedic injuries (for example open | півії | Weuluin | півіі | weuluiti | LOW |
| RH11 | reduction and internal fixation) | Medium | Low | High | Medium | Low |
| RH12 | Repair of cleft lip and cleft palate | Low | Low | Low | Low | Medium |
| RH13 | Repair of club foot | High | Low | Medium | Low | Low |
| RH14 | Cataract extraction | Low | Medium | High | Medium | High |
| RH15 | Repair of anorectal malformations and Hirschsprung's disease | Medium | Low | Low | Low | Low |
| RH16 | Repair of abstetric fistula | Medium | Low | Medium | Low | Low |
| RH17 | Ventriculoperitoneal Shunt | Medium | Low | Low | Low | Low |
| RH18 | Surgery for Trachomatous Trichiasis | Medium | Low | Medium | Medium | Medium |
| RH19 | Referral level hospital pathology services | None | None | None | None | Medium |
| RH20 | Speciality pathology services | None | None | None | None | Medium |
| NELU | ר ארביומוונא אמרוחוחפא אבו אורבא | NOTE | NOTE | NULLE | NOTE | weuluill |